arXiv:math/0411307v1 [math.DG] 13 Nov 2004

HYPER-KÄHLER QUOTIENTS OF SOLVABLE LIE GROUPS

M. L. BARBERIS, I. DOTTI, AND A. FINO

Abstract. In this paper we apply the hyper-Kähler quotient construction to Lie groups with a left invariant hyper-Kähler structure under the action of a closed abelian subgroup by left multiplication. This is motivated by the fact that some known hyper-Kähler metrics can be recovered in this way by considering different Lie group structures on $\mathbb{H}^p \times \mathbb{H}^q$ ($\mathbb{H}$: the quaternions). We obtain new complete hyper-Kähler metrics on Euclidean spaces and give their local expressions.

1. Introduction

Hyper-Kähler reduction allows to construct hyper-Kähler manifolds from others that admit a group acting by tri-holomorphic isometries [8]. Families of $4n$ dimensional hyper-Kähler quotients admitting a tri-holomorphic T^n-action were constructed in [5, 2]. In particular, in [2] the geometry and topology of hyper-Kähler quotients of $\mathbb{H}^d$ by subtori of T^d has been studied.

The hyper-Kähler quotient construction has been also applied in [4] to the flat space $\mathbb{H}^d$ to obtain some monopole moduli space metrics in explicit form using [10, 15], for instance the Taubian-Calabi [16] and the Lee-Weinberg-Yi metric [11]. These are constructed by considering the following actions of $\mathbb{R}$ on $\mathbb{H} \times \mathbb{H}^m$ (resp. $\mathbb{R}^m$ on $\mathbb{H}^m \times \mathbb{H}^m$):

$$\begin{array}{rcl} \mathbb{R} \times \mathbb{H} \times \mathbb{H}^m & \to & \mathbb{H} \times \mathbb{H}^m \\ (t, (q, w_1, \dots, w_m)) & \to & (t + q, e^{it} w_1, \dots, e^{it} w_m), \end{array}$$

$$\begin{array}{rcl} \mathbb{R}^m \times \mathbb{H}^m \times \mathbb{H}^m & \to & \mathbb{H}^m \times \mathbb{H}^m \\ ((t_1, \dots, t_m), (q_1, \dots, q_m, w_1, \dots, w_m)) & \to & (t_1 + q_1, \dots, t_m + q_m, \\ & & e^{i\langle \theta_1, T\rangle} w_1, \dots, e^{i\langle \theta_m, T\rangle} w_m), \end{array}$$

where $\theta \in \mathrm{GL}(m, \mathbb{R})$, $T = (t_1, \dots, t_m)$, θ_β are the rows of θ and $\langle\ ,\ \rangle$ is the Euclidean inner product in $\mathbb{R}^m$. The first action gives rise to the Taubian-Calabi metric, which coincides with the Taub-Nut metric for $q = 1$, and the second one corresponds to the Lee-Weinberg-Yi metric. We show that in both cases the metric can be recovered by endowing $\mathbb{H} \times \mathbb{H}^m$ (resp. $\mathbb{H}^m \times \mathbb{H}^m$) with a hyper-Kähler Lie group structure and taking the quotient with respect to a suitable closed abelian subgroup.

1991 *Mathematics Subject Classification.* 53C26, 22E25, 53D20.

Research partially supported by Conicet, Antorchas, SecytU.N.C (Argentina), by MIUR, CNR (Italy) and ESI (Vienna).

In the present work we study hyper-Kähler quotients starting from a Lie group G with a left invariant hyper-Kähler structure. Such a group is necessarily flat since it is Ricci flat and homogeneous (see [1]). It follows from [12] that G must be 2-step solvable and when G is simply connected, G is a semidirect product of the form $\mathbb{H}^p \ltimes_\theta \mathbb{H}^q$, where θ is a homomorphism from $\mathbb{H}^p$ to T^q, a maximal torus in $\mathrm{Sp}(q)$ (see Proposition 3.1 and (13)). This leads us to get a characterization of hyper-Kähler Lie groups.

We take a connected closed abelian subgroup $\mathbb{R}^l$ ($l \leq p$) of $\mathbb{H}^p$ which acts on G by left translations, hence the action is free and the moment map has no critical points. This action is tri-Hamiltonian, therefore the hyper-Kähler quotient construction [8] can be applied. We prove that the metric obtained on the hyper-Kähler quotient is complete and the quotient is diffeomorphic to an Euclidean space. Since the $\mathbb{R}^l$-action commutes with an action of the torus T^q, if $l = p$ the $4q$-dimensional hyper-Kähler quotient admits a tri-holomorphic T^q-action. Such action has a unique fixed point when $p = q$. In this way we obtain new complete hyper-Kähler metrics which generalize the Taubian-Calabi and the Lee-Weinberg-Yi metrics. Using the same method as in [4, 15, 10], we obtain a local expression of the hyper-Kähler quotient metrics. This expression is given in terms of the structure constants of the corresponding Lie group $\mathbb{H}^p \ltimes_\theta \mathbb{H}^q$.

Acknowledgments. The authors thank Andrew Swann for useful comments and Nigel Hitchin for helpful suggestions. The first and third authors are grateful for the hospitality at the International Erwin Schrödinger Institute for Mathematical Physics in Vienna during the program “Geometric and analytic problems related to Cartan connections ”.

2. Preliminaries

Let $(\mathfrak{g}, g)$ be a metric Lie algebra, that is, $\mathfrak{g}$ is a Lie algebra endowed with an inner product g. The Levi Civita connection associated to the metric can be computed by

$$2g(\nabla_X Y, Z) \;=\; g([X,Y],Z) - g([Y,Z],X) + g([Z,X],Y)), \tag{1}$$

for any X, Y, Z in $\mathfrak{g}$.

A hypercomplex structure on $\mathfrak{g}$ is a triple of complex structures $\{J_\alpha\}_{\alpha=1,2,3}$ satisfying the quaternion relations

$$J_\alpha^2 = -\mathrm{id},\ \alpha = 1,2,3, \qquad J_1 J_2 = -J_2 J_1 = J_3,$$

together with the vanishing of the Nijenhuis tensor $N_\alpha(X,Y) = 0$, for any $X, Y \in \mathfrak{g}$ and $\alpha = 1,2,3$. Here, the Nijenhuis tensor stands for

$$N_\alpha(X,Y) = J_\alpha([X,Y] - [J_\alpha X, J_\alpha Y]) - ([J_\alpha X, Y] + [X, J_\alpha Y]),$$

where $X, Y \in \mathfrak{g}$.

Let $\mathfrak{g}$ be a Lie algebra endowed with a hypercomplex structure $\{J_\alpha\}_{\alpha=1,2,3}$ and an inner product g, compatible with the hypercomplex structure, that is

$$g(X,Y) = g(J_1 X, J_1 Y) = g(J_2 X, J_2 Y) = g(J_3 X, J_3 Y),$$

for all $X, Y \in \mathfrak{g}$. We will say that $(\mathfrak{g}, \{J_\alpha\}, g)$ is a hyper-Kähler Lie algebra when $(\mathfrak{g}, J_\alpha, g)$ is a Kähler Lie algebra, for each α, that is, $\nabla J_\alpha = 0$, where ∇ is the Levi-Civita connection of g. This is equivalent to $d\omega_\alpha = 0$, where ω_α are the associated Kähler forms defined by $\omega_\alpha(X, Y) = g(J_\alpha X, Y),\ X, Y \in \mathfrak{g}$.

If G is a Lie group with Lie algebra $\mathfrak{g}$ then the above structures on $\mathfrak{g}$ can be left translated to all of G obtaining invariant hyper-Kähler structures on G.

Note that a Lie group with an invariant hyper-Kähler structure is necessarily flat since a hyper-Kähler metric is Ricci flat and in the homogeneous case, Ricci flatness implies flatness (see [1]). Examples of non commutative Lie groups carrying a flat invariant metric are given by $T^k \ltimes \mathbb{R}^m$ where T^k is a torus in $\mathrm{SO}(m)$. The next proposition, which is a consequence of the characterization of flat Lie algebras given in [12], shows that this family of examples essentially exhausts the class (see also [3]). This will allow us to give a characterization of hyper-Kähler Lie algebras as a special class of subalgebras of $\mathbb{R}^s \times \mathfrak{e}(4q)$, where $\mathfrak{e}(4q) = \mathfrak{so}(4q) \ltimes \mathbb{R}^{4q}$ is the Euclidean Lie algebra.

Proposition 2.1 ([12]). *Let $(\mathfrak{g}, g)$ be a flat Lie algebra. Then $\mathfrak{g}$ decomposes orthogonally as*

$$\mathfrak{g} = \mathfrak{z}(\mathfrak{g}) \oplus \mathfrak{h} \oplus \mathfrak{g}^1,$$

where $\mathfrak{z}(\mathfrak{g})$ is the center of $\mathfrak{g}$, $\mathfrak{h}$ is an abelian Lie subalgebra, the commutator ideal $\mathfrak{g}^1$ is abelian and the following conditions are satisfied:

i) ad: $\mathfrak{h} \to \mathfrak{so}(\mathfrak{g}^1)$ is injective and $\mathfrak{g}^1$ is even dimensional;

ii) $\mathrm{ad}_X = \nabla_X$ for any $X \in \mathfrak{z}(\mathfrak{g}) \oplus \mathfrak{h}$.

In particular, $\mathfrak{g}$ is isomorphic to a Lie subalgebra of $\mathbb{R}^s \times \mathfrak{e}(\mathfrak{g}^1)$, where $s = \dim \mathfrak{z}(\mathfrak{g})$.

Proof. By [12] a flat Lie algebra $(\mathfrak{g}, g)$ decomposes orthogonally as

$$\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{b}, \tag{2}$$

where $\mathfrak{h}$ is an abelian Lie subalgebra, $\mathfrak{b}$ is the abelian ideal defined by $\{B \in \mathfrak{g} : \nabla_B = 0\}$ and

$$\mathrm{ad}_X : \mathfrak{b} \to \mathfrak{b}$$

is skew-symmetric, for any $X \in \mathfrak{h}$. We observe that the choice of $\mathfrak{b}$, the fact that ∇ is torsion free and that $\mathfrak{h}$ is abelian imply

$$\mathrm{ad}_X = \nabla_X, \qquad \text{for any } X \in \mathfrak{h}. \tag{3}$$

The above equation and the choice of $\mathfrak{b}$ imply

$$\mathrm{ad} : \mathfrak{h} \to \mathfrak{so}(\mathfrak{g}) \tag{4}$$

is injective.

We notice next that the decomposition (2) implies that $\mathfrak{g}^1 \subseteq \mathfrak{b}$, hence $\mathfrak{b}$ decomposes orthogonally as

$$\mathfrak{b} = \mathfrak{v} \oplus \mathfrak{g}^1.$$

We show below that $\mathfrak{v} = \mathfrak{z}(\mathfrak{g})$, where $\mathfrak{z}(\mathfrak{g})$ denotes the center of $\mathfrak{g}$. In particular, $\mathfrak{g}$ will decompose orthogonally as

$$\mathfrak{g} = \mathfrak{z}(\mathfrak{g}) \oplus \mathfrak{h} \oplus \mathfrak{g}^1,$$

with $\mathfrak{h}$ and $\mathfrak{g}^1$ abelian and such that *i*) holds. To show that $\mathfrak{g}^1$ is even dimensional, assume that $\dim \mathfrak{g}^1 = 2m+1$. Since ad_X, $X \in \mathfrak{h}$, is a commutative family of endomorphisms in $\mathfrak{so}(2m+1)$, they are conjugate to elements in a maximal abelian subalgebra of $\mathfrak{so}(2m+1)$, hence there exists $Z \in \mathfrak{g}^1$ such that $\mathrm{ad}_X(Z) = 0$ for any $X \in \mathfrak{h}$, therefore $Z \in \mathfrak{z}(\mathfrak{g}) \cap \mathfrak{g}^1$, a contradiction.

Since $\mathrm{ad}_X : \mathfrak{g}^1 \to \mathfrak{g}^1$ and it is skew-symmetric, for any $X \in \mathfrak{h}$, it preserves $\mathfrak{v}$. Therefore, $[X, \mathfrak{v}] \subset \mathfrak{v} \cap \mathfrak{g}^1 = 0$ for $X \in \mathfrak{h}$ and $\mathfrak{v} \subset \mathfrak{z}(\mathfrak{g})$ follows. On the other hand, if $Y \in \mathfrak{z}(\mathfrak{g})$, then:

$$0 = g([Y, X], U) = g(Y, [X, U]),$$

for every $X \in \mathfrak{h}$, $U \in \mathfrak{g}^1$, that is, $\mathfrak{z}(\mathfrak{g}) \perp \mathfrak{g}^1$ since $\mathfrak{g}^1 = [\mathfrak{h}, \mathfrak{g}^1]$. From $\mathfrak{z}(\mathfrak{g}) \cap \mathfrak{h} = 0$ one has that $\mathfrak{v} = \mathfrak{z}(\mathfrak{g})$.

Finally, using (1) one can compute $\nabla_Y = 0$ for $Y \in \mathfrak{z}(\mathfrak{g})$. This together with (3) imply *ii*) and the proposition follows.

□

We will say that two flat Lie algebras $(\mathfrak{g}_1, g_1)$ and $(\mathfrak{g}_2, g_2)$ are equivalent if there exists an orthogonal Lie algebra isomorphism $\eta : \mathfrak{g}_1 \to \mathfrak{g}_2$. Note that $\eta : \mathfrak{z}(\mathfrak{g}_1) \to \mathfrak{z}(\mathfrak{g}_2)$, $\eta : \mathfrak{g}_1^1 \to \mathfrak{g}_2^1$ and therefore $\eta : \mathfrak{h}_1 \to \mathfrak{h}_2$ (see Proposition 2.1). Let $\mathrm{ad}_i : \mathfrak{h}_i \to \mathfrak{so}(\mathfrak{g}_i^1)$, $i = 1, 2$, be the corresponding injective maps induced by the adjoint representation on $\mathfrak{g}_i$. Then the following diagram is commutative:

$$\begin{tikzcd}
\mathfrak{h}_1 \arrow[r, "\mathrm{ad}_1"] \arrow[d, "\eta"'] & \mathfrak{so}(\mathfrak{g}_1^1) \arrow[d, "I_\eta"] \\
\mathfrak{h}_2 \arrow[r, "\mathrm{ad}_2"] & \mathfrak{so}(\mathfrak{g}_2^1)
\end{tikzcd}$$

where I_η denotes conjugation by η. Conversely, it follows from Proposition 2.1 that every flat Lie algebra with $2m$ dimensional commutator and s dimensional center is equivalent to $\mathbb{R}^s \times \mathbb{R}^k \ltimes_\rho \mathbb{R}^{2m}$, where $\rho : \mathbb{R}^k \to \mathfrak{so}(2m)$ is injective, $\rho(\mathbb{R}^k)\mathbb{R}^{2m} = \mathbb{R}^{2m}$, the only non zero Lie brackets being

$$[X, Y] = \rho(X)Y, \ X \in \mathbb{R}^k, \ Y \in \mathbb{R}^{2m}.$$

Given a flat Lie algebra $\mathbb{R}^s \times (\mathbb{R}^k \ltimes_\rho \mathbb{R}^{2m})$, the family $\{\rho(T) : T \in \mathbb{R}^k\} \subseteq \mathfrak{so}(2m)$ is an abelian subalgebra, then it is conjugate by an element in $SO(2m)$ to a subalgebra of the following maximal abelian subalgebra of $\mathfrak{so}(2m)$:

$$\mathfrak{t}^m = \left\{ \begin{pmatrix} 0 & -\phi_1 & & & \\ \phi_1 & 0 & & & \\ & & \ddots & & \\ & & & 0 & -\phi_m \\ & & & \phi_m & 0 \end{pmatrix} : \phi_\alpha \in \mathbb{R} \right\}$$

with respect to an orthonormal basis $\{f_1,\dots,f_{2m}\}$ of $\mathbb{R}^{2m}$. In particular, $k\le m$ and we may assume that any flat Lie algebra is equivalent to a Lie algebra such that $\rho(\mathbb{R}^k)\subset\mathfrak{t}^m$.

Let $\theta=(\theta^\alpha_\beta)$ be the real $m\times k$ matrix of rank k such that

$$\rho(e_\alpha)=\begin{pmatrix} 0 & -\theta^\alpha_1 & & & \\ \theta^\alpha_1 & 0 & & & \\ & & \ddots & & \\ & & & 0 & -\theta^\alpha_m \\ & & & \theta^\alpha_m & 0 \end{pmatrix},\quad 1\le\alpha\le k, \tag{5}$$

where $\{e_1,\dots,e_k\}$ is an orthonormal basis of $\mathbb{R}^k$. The condition $\rho(\mathbb{R}^k)\mathbb{R}^{2m}=\mathbb{R}^{2m}$ is equivalent to the fact that every row θ_β of θ is non zero.

We introduce some notation that will be used in the next result. Let $M(k,m;k)$ be the set of $m\times k$ real matrices of rank k. $M(k,m;k)$ can be viewed inside $\operatorname{End}(\mathbb{R}^k,\mathfrak{so}(2m))$ by means of the inclusion ρ:

$$M(k,m;k)\hookrightarrow\operatorname{End}(\mathbb{R}^k,\mathfrak{so}(2m)),\qquad \theta\mapsto\rho_\theta.$$

We identify $M(k,m;k)$ with its image under ρ and let $O(k)\times O(2m)$ act on $M(k,m;k)$ as follows:

$$O(k)\times O(2m)\times M(k,m;k)\to M(k,m;k),\qquad (A,B,\ \rho_\theta)\mapsto B\rho_{(\theta A)}B^{-1}, \tag{6}$$

where $B\rho_\theta B^{-1}\in\operatorname{End}(\mathbb{R}^k,\mathfrak{so}(2m))$ is defined by $B\rho_\theta B^{-1}(T)=B\,\rho_\theta(T)\,B^{-1}$, $T\in\mathbb{R}^k$. It follows from the definition of equivalence between flat Lie algebras that

$$\mathbb{R}^k\ltimes_{\rho_\theta}\mathbb{R}^{2m}\cong\mathbb{R}^k\ltimes_{\rho_{\theta'}}\mathbb{R}^{2m}$$

if and only if ρ_θ and $\rho_{\theta'}$ lie in the same $O(k)\times O(2m)$-orbit.

The next proposition summarizes the above results and gives the classification of flat Lie algebras that will be needed in the next section (see also [9]).

Proposition 2.2. *Let $(\mathfrak{g},g)$ be a flat Lie algebra, $\dim\mathfrak{g}^1=2m$, $\dim\mathfrak{z}(\mathfrak{g})=s$. Then there exists $\theta=(\theta^\alpha_\beta)\in M(k,m;k)$ such that $\theta_\beta\ne0$ for every $1\le\beta\le m$ and $\mathfrak{g}$ decomposes orthogonally as*

$$\mathfrak{g}\cong\mathbb{R}^s\times(\mathbb{R}^k\ltimes_{\rho_\theta}\mathbb{R}^{2m}),$$

where $\mathbb{R}^k\ltimes_{\rho_\theta}\mathbb{R}^{2m}$ has an orthonormal basis $\{e_1,\dots,e_k,f_1,\dots,f_{2m}\}$ and $T\in\mathbb{R}^k$ acts on $\mathbb{R}^{2m}$ in the following way:

$$\rho_\theta(T)=\begin{pmatrix} 0 & -\langle T,\theta_1\rangle & & & \\ \langle T,\theta_1\rangle & 0 & & & \\ & & \ddots & & \\ & & & 0 & -\langle T,\theta_m\rangle \\ & & & \langle T,\theta_m\rangle & 0 \end{pmatrix} \tag{7}$$

where $\langle\ ,\ \rangle$ denotes the Euclidean inner product on $\mathbb{R}^k$. Moreover,

$$\mathbb{R}^k\ltimes_{\rho_\theta}\mathbb{R}^{2m}\cong\mathbb{R}^k\ltimes_{\rho_{\theta'}}\mathbb{R}^{2m}$$

as flat Lie algebras if and only if ρ_θ *and* $\rho_{\theta'}$ *lie in the same* $O(k) \times O(2m)$*-orbit under the action* (6).

Remark. Note that the Lie algebra $\mathbb{R}^k \ltimes_{\rho_\theta} \mathbb{R}^{2m}$ is a Lie subalgebra of the Euclidean Lie algebra $\mathfrak{e}(2m)$:

$$\mathbb{R}^k \ltimes_{\rho_\theta} \mathbb{R}^{2m} \hookrightarrow \mathfrak{e}(2m), \quad (T, W) \mapsto \begin{pmatrix} \rho_\theta(T) & W \\ 0 & 0 \end{pmatrix},$$

$T \in \mathbb{R}^k$, $W \in \mathbb{R}^{2m}$. However, the inner product on $\mathbb{R}^k \ltimes_{\rho_\theta} \mathbb{R}^{2m}$ is not the one induced from $\mathfrak{e}(2m)$.

The next corollary follows from the description given in Proposition 2.2.

Corollary 2.1. *Any even dimensional flat Lie algebra is Kähler flat.*

Proof. Let $\mathfrak{g}_\theta = \mathbb{R}^s \times (\mathbb{R}^k \ltimes_{\rho_\theta} \mathbb{R}^{2m})$ be as in Proposition 2.2 and J the endomorphism of $\mathfrak{g}_\theta$ such that $J^2 = -\mathrm{id}$, $Jf_{2i+1} = f_{2i}$, $i = 0, \dots, m-1$, and J is orthogonal preserving $\mathbb{R}^s \times \mathbb{R}^k$. The integrability of J, that is, the vanishing of N_J, follows from $\rho_\theta(T)J = J\rho_\theta(T)$, for any $T \in \mathbb{R}^k$. Moreover, $\nabla J = 0$ since $\nabla_T = \rho_\theta(T)$, for $T \in \mathbb{R}^k$. Therefore $(\mathfrak{g}, J, g)$ is Kähler flat. □

3. Hyper-Kähler Lie groups

We apply Proposition 2.1 to give a characterization of the Lie algebras carrying a hyper-Kähler structure $(\{J_\alpha\}, g)$.

Proposition 3.1. *Let* $(\mathfrak{g}, \{J_\alpha\}, g)$*,* $\alpha = 1, 2, 3$*, be a hyper-Kähler Lie algebra. Then* $\mathfrak{g}$ *decomposes orthogonally as*

$$\mathfrak{g} = \mathfrak{t} \oplus \mathfrak{g}^1, \qquad \mathfrak{z}(\mathfrak{g}) \subset \mathfrak{t},$$

with both $\mathfrak{t}$ *and* $\mathfrak{g}^1$ *abelian and* J_α*-invariant,* $\alpha = 1, 2, 3$*, such that*
i) $ad_X J_\alpha = J_\alpha ad_X$*, for any* $X \in \mathfrak{t}$*,* $\alpha = 1, 2, 3$*;*
ii) $g(ad_X Y, Z) + g(Y, ad_X Z) = 0$*, for any* $X \in \mathfrak{t}, Y, Z \in \mathfrak{g}$*.*

Proof. Since a hyper-Kähler Lie algebra is flat [1], $\mathfrak{g}$ decomposes orthogonally as $\mathfrak{g} = \mathfrak{z}(\mathfrak{g}) \oplus \mathfrak{h} \oplus \mathfrak{g}^1$ and the conditions of Proposition 2.1 are satisfied. Set

$$\mathfrak{t} = \mathfrak{z}(\mathfrak{g}) \oplus \mathfrak{h}.$$

We show next that if $(\mathfrak{g}, \{J_\alpha\}, g)$, $\alpha = 1, 2, 3$, is a hyper-Kähler Lie algebra then $\mathfrak{t}$ and $\mathfrak{g}^1$ are J_α-invariant, $\alpha = 1, 2, 3$, and that condition *i*) is satisfied. Observe that if $X \in \mathfrak{t}$ and $B \in \mathfrak{g}^1$, using that $\nabla J_\alpha = 0$ and *ii*) of Proposition 2.1, one has

$$J_\alpha[X, B] = J_\alpha \nabla_X B = [X, J_\alpha B],$$

therefore, *i*) follows. Since $\mathfrak{g}^1 = [\mathfrak{h}, \mathfrak{g}^1]$, the above equation also implies that $\mathfrak{g}^1$ is J_α-invariant and the decomposition $\mathfrak{t} \oplus \mathfrak{g}^1$ satisfies the desired properties.

□

We will say that two hyper-Kähler Lie algebras $(\mathfrak{g}, \{J_\alpha\}, g)$ and $(\mathfrak{g}', \{J'_\alpha\}, g')$ are equivalent if there exists an equivalence η of metric Lie algebras such that $\eta J_\alpha = J'_\alpha \eta$, $\alpha = 1, 2, 3$.

Consider the hypercomplex structure on

$$\mathbb{H}^q = \{(W_1, \dots, W_q) : W_\alpha = u_\alpha + y_\alpha i + z_\alpha j + w_\alpha k \ : \ u_\alpha, y_\alpha, z_\alpha, w_\alpha \in \mathbb{R}\}$$

given by right multiplication by $-i$, $-j$, $-k$:

$$J_1 = R_{-i}, \qquad J_2 = R_{-j}, \qquad J_3 = R_{-k}.$$

We identify $\mathbb{H}^q \cong \mathbb{R}^{4q}$ with the Euclidean metric and let $\mathrm{Sp}(q) = O(4q) \cap \mathrm{GL}\,(q, \mathbb{H})$, where

$$\mathrm{GL}(q, \mathbb{H}) = \{T \in \mathrm{GL}(4q, \mathbb{R}) : TJ_\alpha = J_\alpha T, \ \alpha = 1, 2, 3\}.$$

Let $\mathfrak{t}^q$ be the following maximal abelian subalgebra of the Lie algebra $\mathfrak{sp}(q)$ of $\mathrm{Sp}(q)$:

$$\mathfrak{t}^q = \left\{ \begin{pmatrix} 0 & -\phi_1 & 0 & 0 & & & & & \\ \phi_1 & 0 & 0 & 0 & & & & & \\ 0 & 0 & 0 & -\phi_1 & & & & & \\ 0 & 0 & \phi_1 & 0 & & & & & \\ & & & & \ddots & & & & \\ & & & & & 0 & -\phi_q & 0 & 0 \\ & & & & & \phi_q & 0 & 0 & 0 \\ & & & & & 0 & 0 & 0 & -\phi_q \\ & & & & & 0 & 0 & \phi_q & 0 \end{pmatrix} \ : \ \phi_i \in \mathbb{R} \right\}. \tag{8}$$

We obtain the analogue of Proposition 2.2 using the same procedure as before. Observe that, in this case, $\mathbb{R}^s \times (\mathbb{R}^k \ltimes_{\rho_\theta} \mathbb{H}^q) \cong \mathbb{R}^s \times (\mathbb{R}^k \ltimes_{\rho_{\theta'}} \mathbb{H}^q)$ as hyper-Kähler Lie algebras if and only if ρ_θ and $\rho_{\theta'}$ lie in the same $\mathrm{Sp}(p) \times \mathrm{Sp}(q)$-orbit, where $s + k = 4p$ and the action of $\mathrm{Sp}(p) \times \mathrm{Sp}(q)$ is the analogue of (6).

Proposition 3.2. *Let $(\mathfrak{g}, \{J_\alpha\}, g)$ be a hyper-Kähler Lie algebra with $\dim \mathfrak{g}^1 = 4q$ and $\dim \mathfrak{z}(\mathfrak{g}) = s$. Then there exists $\theta = (\theta^\alpha_\beta) \in M(k, q; k)$, $s + k = 4p$, such that $\theta_\beta \neq 0$ for $1 \leq \beta \leq q$ and*

$$\mathfrak{g} \cong \mathbb{R}^s \times (\mathbb{R}^k \ltimes_{\rho_\theta} \mathbb{H}^q).$$

$\mathbb{R}^k \ltimes_{\rho_\theta} \mathbb{H}^q$ is the Lie algebra with orthonormal basis

$$\{e_1, \dots, e_k, f_1, f_1 i, f_1 j, f_1 k, \dots, f_q, f_q i, f_q j, f_q k\}$$

such that an element $T \in \mathbb{R}^k$ acts on $\mathbb{H}^q$ by

$$\rho_\theta(T) = \begin{pmatrix} \rho^1_\theta(T) & & \\ & \ddots & \\ & & \rho^q_\theta(T) \end{pmatrix}, \tag{9}$$

where $\rho^\beta_\theta(T)$ *is the following* 4×4 *real matrix:*

$$\rho^\beta_\theta(T) = \begin{pmatrix} 0 & -\langle T,\theta_\beta\rangle & 0 & 0 \\ \langle T,\theta_\beta\rangle & 0 & 0 & 0 \\ 0 & 0 & 0 & -\langle T,\theta_\beta\rangle \\ 0 & 0 & \langle T,\theta_\beta\rangle & 0 \end{pmatrix}$$

and $\langle\ ,\ \rangle$ *denotes the Euclidean inner product on* $\mathbb{R}^k$. *The Lie algebra* $\mathbb{R}^s\times(\mathbb{R}^k\ltimes_{\rho_\theta}\mathbb{H}^q)$ *is hyper-Kähler with its natural hypercomplex structure obtained by extending* $R_{-i},\ R_{-j},\ R_{-k}$ *on* $\mathbb{H}^q$ *with any pair of anticommuting complex endomorphisms on* $\mathbb{R}^s\times\mathbb{R}^k$ *and the canonical inner product. Moreover,*

$$\mathbb{R}^s\times(\mathbb{R}^k\ltimes_{\rho_\theta}\mathbb{H}^q)\cong\mathbb{R}^s\times(\mathbb{R}^k\ltimes_{\rho_{\theta'}}\mathbb{H}^q)$$

as hyper-Kähler Lie algebras if and only if ρ_θ *and* $\rho_{\theta'}$ *lie in the same* $Sp(p)\times Sp(q)$*-orbit.*

3.1. **Examples.** As a consequence of Proposition 3.2 we have that there is a one parameter family of 8-dimensional hyper-Kähler Lie algebras $\mathfrak{g}_\theta$:

$$\mathfrak{g}_\theta\cong\mathbb{R}^3\times(\mathbb{R}\ltimes_\theta\mathbb{H}), \tag{10}$$

where $\mathbb{R}\ltimes_\theta\mathbb{H}$ has an orthonormal basis $\{e_1, f_1, f_1 i, f_1 j, f_1 k\}$ and e_1 acts on $\mathbb{H}$ as follows:

$$\rho_\theta(e_1) = \begin{pmatrix} 0 & -\theta & & \\ \theta & 0 & & \\ & & 0 & -\theta \\ & & \theta & 0 \end{pmatrix}.$$

Note that these are pairwise non equivalent flat metric Lie algebras for different values of θ, but they are isomorphic Lie algebras for $\theta\neq 0$.

In dimension 12 there are infinitely many non isomorphic Lie algebra structures admitting hyper-Kähler metrics. In fact, for a fixed real number $s\neq 0$ we define $\mathfrak{g}_s = \mathbb{R}^3\times(\mathbb{R}\ltimes_s\mathbb{H}^2)$, where $\mathbb{R}\ltimes_s\mathbb{H}^2$ has an orthonormal basis as in the statement of Proposition 3.2 with e_1 acting on $\mathbb{H}^2$ as follows:

$$\rho_s(e_1) = \begin{pmatrix} 0 & -1 & & & & & & \\ 1 & 0 & & & & & & \\ & & 0 & -1 & & & & \\ & & 1 & 0 & & & & \\ & & & & 0 & -s & & \\ & & & & s & 0 & & \\ & & & & & & 0 & -s \\ & & & & & & s & 0 \end{pmatrix}.$$

It turns out that $\mathfrak{g}_s$ and $\mathfrak{g}_r$ are non isomorphic for $s\neq r$.

We describe below the Lie bracket on $\mathfrak{g}_\theta = \mathbb{R}^k \ltimes_{\rho_\theta} \mathbb{H}^q$:

$$\begin{aligned}[(X,W),(X',W')] =&(0, i(\langle X,\theta_1\rangle W_1' - \langle X',\theta_1\rangle W_1), \dots, i(\langle X,\theta_q\rangle W_q' - \langle X',\theta_q\rangle W_q)) \\ =&(0, \rho_\theta(X)W' - \rho_\theta(X')W),\end{aligned} \tag{11}$$

$X, X' \in \mathbb{R}^k$.

The product on the simply connected Lie group $G_\theta = \mathbb{R}^k \ltimes_\theta \mathbb{H}^q$ with Lie algebra $\mathfrak{g}_\theta$ is given as follows:

$$(X,W)\cdot(X',W') = (X+X',\, W + \theta(X)W'), \tag{12}$$

where $X, X' \in \mathbb{R}^k,\ W, W' \in \mathbb{H}^q,\ W' = (W_1', \dots, W_q')$ and

$$\theta(X)W' = (e^{i\langle X,\theta_1\rangle}W_1', \dots, e^{i\langle X,\theta_q\rangle}W_q'). \tag{13}$$

Using that $(X,W)^{-1} = -(X, \theta(-X)W)$, conjugation by (X,W) is given as follows:

$$I_{(X,W)}(X',W') = (X,W)\cdot(X',W')\cdot(X,W)^{-1} = (X', W + \theta(X)W' - \theta(X')W)$$

and therefore

$$\mathrm{Ad}(X,W)(X',W') = (X', \theta(X)W') + [(0,W),(X',0)] = (X', \theta(X)W' - \rho_\theta(X')W), \tag{14}$$

for $X, X' \in \mathbb{R}^k$, $W, W' \in \mathbb{H}^q$.

The left invariant flat metric g on $\mathbb{R}^k \ltimes_{\rho_\theta} \mathbb{H}^q$ in coordinates $(x_1, \dots, x_k, W_1, \dots, W_q)$, where $W_j = (u_j, y_j, z_j, w_j)$, is the Euclidean metric

$$g = \sum_{j=1}^{k} dx_j^2 + \sum_{j=1}^{q}(du_j^2 + dy_j^2 + dz_j^2 + dw_j^2).$$

We will need to express the Euclidean metric on $\mathbb{H}^q$ in coordinates adapted to the hyper-Kähler moment map. Any quaternion may be written as

$$W_\beta = e^{i\psi_\beta/2}a_\beta, \quad \beta = 1, \dots, q,$$

with $\psi_\beta \in (0, 4\pi]$ and a_β is pure imaginary, so that $\overline{a}_\beta = -a_\beta$. Let

$$\mathbf{r}_\beta = \overline{W}_\beta i W_\beta = \overline{a}_\beta i a_\beta = -a_\beta i a_\beta.$$

The flat metric on $\mathbb{H}^q$ in coordinates $(\psi_\beta, \mathbf{r}_\beta)$, $\beta = 1, \dots, q$, is given by

$$\frac{1}{4}\sum_{\beta=1}^{q}\left(\frac{1}{r_\beta}d\mathbf{r}_\beta^2 + r_\beta(d\psi_\beta + \mathbf{\Omega}_\beta \cdot d\mathbf{r}_\beta)^2\right), \tag{15}$$

where

$$r_\beta = |\mathbf{r}_\beta|, \quad \mathrm{curl}(\mathbf{\Omega}_\beta) = \mathrm{grad}\left(\frac{1}{r_\beta}\right)$$

(the curl and grad operations are taken with respect to the Euclidean metric on $\mathbb{R}^3$ with cartesian coordinates $\mathbf{r}_\beta$).

4. Completeness of the hyper-Kähler quotient metric

According to Proposition 3.2 any simply connected Lie group with a left invariant hyper-Kähler structure is of the form $G_\theta = \mathbb{R}^s \times (\mathbb{R}^k \ltimes_\theta \mathbb{H}^q)$ ($k \leq q$, $s + k = 4p$) with the hyper-Kähler metric $g = g_1 \times g_2$, where g_1 is the Euclidean metric on $\mathbb{R}^s \times \mathbb{R}^k$ and g_2 is the Euclidean metric on $\mathbb{H}^q$. Let $\mathfrak{g}_\theta$ be the Lie algebra of G_θ. The associated Kähler forms:

$$\omega_\alpha((X_1, W_1), (X_2, W_2)) = g(J_\alpha(X_1, W_1), (X_2, W_2)), \qquad (X_1, W_1), (X_2, W_2) \in \mathfrak{g}_\theta,$$

$\alpha = 1, 2, 3$, when left translated to G_θ become:

$$\omega_\alpha = \omega^1_\alpha + \omega^2_\alpha,$$

where ω^j_α, $j = 1, 2$, $\alpha = 1, 2, 3$ are the standard symplectic forms on a vector space. Therefore, $(G_\theta, g, \omega_\alpha)$ is equivalent, as a hyper-Kähler manifold, to the product

$$(\mathbb{R}^s \times \mathbb{R}^k, g_1, \{\omega^1_\alpha\}) \times (\mathbb{H}^q, g_2, \{\omega^2_\alpha\}). \tag{16}$$

We will apply the hyper-Kähler quotient construction due to [8] to the case when L is the connected closed abelian Lie subgroup $\mathbb{R}^l \subset \mathbb{R}^k$ with Lie algebra $\mathfrak{l} = \mathrm{span}_{\mathbb{R}}\{e_1, \ldots, e_l\}$ such that $\mathfrak{l}$ is isotropic with respect to ω_α for each α. The action of L on G_θ will be given by left translations, therefore it preserves the hyper-Kähler structure. We recall next the quotient construction in our particular case.

Let $\mathcal{X}_V$ be the vector field generated by the action of L, that is, the right invariant vector field such that $\mathcal{X}_{V_e} = V$, where $V \in \mathfrak{l}$. Observe that

$$0 = L_{\mathcal{X}_V}\omega_\alpha = d(i(\mathcal{X}_V)\omega_\alpha) + i(\mathcal{X}_V)d\omega_\alpha,$$

where $i(\mathcal{X}_V)\omega_\alpha$ denotes the 1-form obtained by taking the interior product with $\mathcal{X}_V$. Since the action is symplectic with respect to ω_α, $\alpha = 1, 2, 3$, we have that $i(\mathcal{X}_V)\omega_\alpha$, $\alpha = 1, 2, 3$, is closed. G_θ is simply connected, thus $H^1_{dR}(G_\theta, \mathbb{R}) = \{0\}$ and $i(\mathcal{X}_V)\omega_\alpha$ is exact, that is,

$$i(\mathcal{X}_V)\omega_\alpha = d\,{\mu^{\theta}_\alpha}^V,$$

where ${\mu^{\theta}_\alpha}^V$ is a Hamiltonian function associated to V. Putting all these functions together, we obtain a map to the dual space of the Lie algebra of L

$${\mu^\theta}_\alpha : G_\theta \to \mathfrak{l}^*$$

defined by

$${\mu^\theta}_\alpha(X, W)(V) = {\mu^\theta}^V_\alpha(X, W).$$

L acts on $\mathfrak{l}^*$ by the coadjoint action. When the ambiguities in the choices of ${\mu^{\theta}_\alpha}^V$ may be adjusted to make μ^θ_α L-equivariant, one has the hyper-Kähler moment map

$$\mu^\theta : G_\theta \to \mathfrak{l}^* \otimes \mathrm{Im}\,\mathbb{H},$$

defined by $\mu^\theta = \mu_1^\theta i + \mu_2^\theta j + \mu_3^\theta k$. Our choice of L implies that μ_α^θ is L-equivariant for each α. Indeed, the action A of L on G_θ given by left translations:

$$\begin{array}{rcl} A : L \times G_\theta & \to & G_\theta \\ ((V,0),(X,W)) & \to & (V,0)\cdot(X,W) = (V+X,\, \theta(V)W) \end{array} \tag{17}$$

(recall (12)) can be viewed as a diagonal action of L:

$$A(V)(X,W) = (A_1(V)X, A_2(V)W),$$

where A_1 acts by left translations on $\mathbb{R}^s \times \mathbb{R}^k$ and A_2 is a linear symplectic action on $\mathbb{H}^q$. The moment map μ_α^θ corresponding to A can be obtained by adding up the moment maps of A_1 and A_2 since (16) holds. By a direct calculation one has (see [6]):

$$\mu_\alpha^\theta(X,W)(V) = \omega_\alpha(V,X) + \frac{1}{2}\omega_\alpha(\rho_\theta(V)W,W)$$

(see (11)). The L-equivariance of the first term follows since L is isotropic and the second term is L-equivariant since it is the moment map of a linear action on a symplectic vector space (see [6]).

Let $\xi \in \mathfrak{z}\otimes\mathbb{R}^3$ be a regular value for μ^θ, where $\mathfrak{z}$ is the subspace of $\mathfrak{l}^*$ of invariant elements under the coadjoint action, and consider the quotient $L\backslash(\mu^\theta)^{-1}(\xi)$. According to [8], when the action of L on $(\mu^\theta)^{-1}(\xi)$ is free with Hausdorff quotient manifold $L\backslash(\mu^\theta)^{-1}(\xi)$, the hyper-Kähler metric on G_θ induces a hyper-Kähler metric on $L\backslash(\mu^\theta)^{-1}(\xi)$.

Our hypotheses imply:

1. the center of $\mathfrak{l}^*$, that is, the invariant elements under the coadjoint action, coincides with $\mathfrak{l}^*$ since L is abelian;
2. the action of L on G_θ is free, hence it is free on $(\mu^\theta)^{-1}(\xi)$, for any $\xi = \xi_1 i + \xi_2 j + \xi_3 k \in \operatorname{Im}\mu^\theta$. In particular, any $\xi \in \operatorname{Im}\mu^\theta$ is a regular value of the hyper-Kähler moment map;
3. since L is closed in G_θ and acts by left translations, the set of right cosets $L\backslash G_\theta$ is a Hausdorff manifold and so is $L\backslash(\mu^\theta)^{-1}(\xi)$, for any $\xi \in \operatorname{Im}\mu^\theta$.

Therefore, $L\backslash(\mu^\theta)^{-1}(\xi)$ inherits a hyper-Kähler metric.

We show next that the hyper-Kähler metric on $L\backslash(\mu^\theta)^{-1}(\xi)$ is always complete. The left invariant metric g on G_θ induces in a natural way a metric $\tilde{g}$ on $L\backslash G_\theta$ such that the natural projection

$$\pi : (G_\theta, g) \to (L\backslash G_\theta, \tilde{g})$$

is a Riemannian submersion. The completeness of g implies that $\tilde{g}$ is also complete (see [7]). Since ξ is a regular value of μ^θ, $(\mu^\theta)^{-1}(\xi)$ is a closed embedded submanifold of G_θ and one has the following commutative diagram:

$$\begin{tikzcd} G_\theta \arrow[r, "\pi"] & L\backslash G_\theta \\ {\mu^\theta}^{-1}(\xi) \arrow[u] \arrow[r, "\tilde{\pi}"] & L\backslash {\mu^\theta}^{-1}(\xi), \arrow[u] \end{tikzcd}$$

where the vertical arrows are the natural inclusions.

Since μ^θ is L-equivariant, it induces a map

$$\tilde{\mu}^\theta : L\backslash G_\theta \to \mathfrak{l}^* \otimes \mathrm{Im}\mathbb{H}$$

and $L\backslash {\mu^\theta}^{-1}(\xi) = \tilde{\mu^\theta}^{-1}(\xi)$. Then $L\backslash {\mu^\theta}^{-1}(\xi)$ is a closed d-dimensional embedded submanifold of $L\backslash G_\theta$ ($d = \dim G_\theta - 4 \dim L$) and the induced metric is complete. By O'Neill formula ([14, 3, Corollary 1]) the sectional curvature of $L\backslash G_\theta$ is non negative. Note that $\mathfrak{l} \subset [\mathfrak{g}_\theta, \mathfrak{g}_\theta]^\perp$, hence the fibers of π are totally geodesic since $\nabla_T V = 0$ for $T, V \in \mathfrak{l}$ (see Proposition 2.1).

We summarize the above paragraphs as follows:

Theorem 4.1. *Let $(G_\theta, \{J_\alpha\}, g)$ be a simply connected hyper-Kähler Lie group, so that $G_\theta = \mathbb{R}^s \times (\mathbb{R}^k \ltimes_\theta \mathbb{H}^q)$, $k \leq q$, $s + k = 4p$. Fix a connected closed abelian isotropic subgroup $L \subset \mathbb{R}^k$ acting on G_θ by the action A as in* (17) *and denote by $\pi : G_\theta \to L\backslash G_\theta$ the associated Riemannian submersion. Then*

1. *the action A of L on G_θ is free and preserves both, the metric g and the symplectic forms ω_α, $\alpha = 1, 2, 3$. The L-equivariant moment map is $\mu^\theta = \mu_1^\theta i + \mu_2^\theta j + \mu_3^\theta k$, with μ_α^θ given by*

$$\mu_\alpha^\theta(X, W)(V) = \omega_\alpha(V, X) + \frac{1}{2}\omega_\alpha(\rho_\theta(V)W, W),$$

   for any $X \in \mathbb{R}^s \times \mathbb{R}^k, W \in \mathbb{H}^q, V \in \mathfrak{l}$;
2. *$L\backslash G_\theta$ has non negative sectional curvature. Moreover, the fibers of π are totally geodesic;*
3. *for any $\xi \in Im\, \mu^\theta$, $L\backslash {\mu^\theta}^{-1}(\xi)$ is a closed embedded submanifold of $L\backslash G_\theta$ and the hyper-Kähler metric on the quotient $L\backslash(\mu^\theta)^{-1}(\xi)$ is complete.*

5. Examples

In the next examples we show that it is possible to describe families of known hyper-Kähler metrics [4] in a unified way, by applying the quotient construction to hyper-Kähler Lie groups G with the action of a suitable abelian subgroup L by left translations.

5.1. **Taub-Nut metric.** Let $\mathfrak{g}_\theta$ be the one parameter family of hyper-Kähler Lie algebras in dimension 8 (see (10)) and G_θ the corresponding simply connected Lie groups. Let $\mathbb{R}$ be the subgroup of $G_\theta = \mathbb{H} \ltimes_\theta \mathbb{H}$ given by $(t, 0)$, $t \in \mathbb{R}$, and let it act on G_θ by left translations, that is:

$$\begin{array}{ccc} \mathbb{R} \times G_\theta & \to & G_\theta \\ (t, (q, w)) & \to & (t + q, e^{i\theta t} w). \end{array}$$

Observe that $\operatorname{Im}\mathbb{H}$ acts trivially on the second factor. The corresponding hyper-Kähler moment map is

$$\begin{aligned}\mu^\theta &= -\operatorname{Im}(q) - \frac{\theta}{2}\left(\operatorname{Re}(iwi\overline{w})i + \operatorname{Re}(iwj\overline{w})j + \operatorname{Re}(iwk\overline{w})k\right)\\ &= -\operatorname{Im}(q) + \frac{\theta}{2}\overline{w}iw.\end{aligned}$$

It can be checked that μ^θ is L-equivariant. The complete hyper-Kähler metric on $\mathbb{R}\backslash(\mu^\theta)^{-1}(0)$ is the Taub-Nut metric with parameter θ^{-1} [4].

5.2. Generalized Taubian-Calabi metric.

Let $\theta = (\theta_1, \dots, \theta_m) \in \mathbb{R}^m$, $G_\theta = \mathbb{H} \ltimes_\theta \mathbb{H}^m$ and $\mathbb{R} = \{(t, 0) : t \in \mathbb{R}\}$ acting on G_θ by left translations:

$$\begin{aligned}\mathbb{R} \times G_\theta &\to G_\theta\\ (t, (q, w_1, \dots, w_m)) &\to (t + q, e^{i\theta_1 t}w_1, \dots, e^{i\theta_m t}w_m).\end{aligned}$$

For $m = 1$ this is the Lie group considered in the first example. The corresponding hyper-Kähler moment map is

$$\begin{aligned}\mu^\theta &= -\operatorname{Im}(q) - \frac{1}{2}\sum_{\beta=1}^{m}\theta_\beta(\operatorname{Re}(iw_\beta i\overline{w_\beta})i + \operatorname{Re}(iw_\beta j\overline{w_\beta})j + \operatorname{Re}(iw_\beta k\overline{w_\beta})k)\\ &= -\operatorname{Im}(q) + \frac{1}{2}\sum_{\beta=1}^{m}\theta_\beta\overline{w_\beta}iw_\beta.\end{aligned}$$

When $\theta_\beta = 1$, for each β, the complete hyper-Kähler metric on $\mathbb{R}\backslash(\mu^\theta)^{-1}(0)$ coincides with the Taubian-Calabi metric [16, 4].

5.3. Lee-Weinberg-Yi metric.

Let $\theta \in \mathrm{GL}(m, \mathbb{R})$, $G_\theta = \mathbb{H}^m \ltimes_\theta \mathbb{H}^m$ and

$$\mathbb{R}^m = \{((t_1, \dots, t_m), 0) : t_i \in \mathbb{R}\}$$

acting on G_θ by left translations:

$$\begin{aligned}\mathbb{R}^m \times G_\theta &\to G_\theta\\ ((t_1, \dots, t_m), (q_1, \dots, q_m, w_1, \dots, w_m)) &\to (t_1 + q_1, \dots, t_m + q_m,\\ &\qquad e^{i\langle\theta_1, T\rangle}w_1, \dots, e^{i\langle\theta_m, T\rangle}w_m),\end{aligned}$$

where $T = (t_1, \dots, t_m)$, θ_β are the rows of θ and $\langle\ ,\ \rangle$ is the Euclidean inner product in $\mathbb{R}^m$. The corresponding hyper-Kähler moment map is

$$\mu^\theta = \left(-\operatorname{Im}(q_1) + \frac{1}{2}\sum_{\beta=1}^{m}\theta^1_\beta\overline{w_\beta}iw_\beta, \dots, -\operatorname{Im}(q_m) + \frac{1}{2}\sum_{\beta=1}^{m}\theta^m_\beta\overline{w_\beta}iw_\beta\right).$$

The complete hyper-Kähler metric on $\mathbb{R}\backslash\mu_\theta^{-1}(0)$ is the Lee-Weinberg-Yi metric with $(\lambda^a_b) = \theta^{-1}$ [11, 13, 4].

6. Topology of the quotient and local description of the metric

Let G_θ be the simply connected hyper-Kähler Lie group $\mathbb{R}^s \times (\mathbb{R}^k \ltimes_\theta \mathbb{H}^q)$ ($s+k=4p$, $k \le q$), $\theta \in M(k,q;k)$ and $\theta(\mathbb{H}^q)=\mathbb{H}^q$. Let $L \subset \mathbb{R}^k$ be a closed abelian subgroup with Lie algebra $\mathfrak{l} = \mathrm{span}_{\mathbb{R}}\{e_1,\dots,e_l\}$ such that $\mathfrak{l}$ is isotropic with respect to ω_α, for each α.

Let T^q be the maximal torus of $\mathrm{Sp}(q)$ with Lie algebra $\mathfrak{t}^q$ (defined in (8)) whose elements are of the form:

$$B = \begin{pmatrix} B(\phi_1) & & \\ & \ddots & \\ & & B(\phi_q) \end{pmatrix}, \tag{18}$$

where $\phi_\beta \in \mathbb{R}$ and $B(\phi_\beta)$ is the following 4×4 real matrix:

$$\begin{pmatrix} \cos(\phi_\beta) & -\sin(\phi_\beta) & 0 & 0 \\ \sin(\phi_\beta) & \cos(\phi_\beta) & 0 & 0 \\ 0 & 0 & \cos(\phi_\beta) & -\sin(\phi_\beta) \\ 0 & 0 & \sin(\phi_\beta) & \cos(\phi_\beta) \end{pmatrix}.$$

We have an action φ of T^q on G_θ:

$$\begin{aligned} &\varphi : T^q \times G_\theta \to G_\theta \\ &(g,(X,W)) \to \varphi(g,(X,W)) = (X, BW), \end{aligned} \tag{19}$$

where BW stands for the product of the $4q\times 4q$ matrix B given in (18) by the column vector $W \in \mathbb{H}^q \cong \mathbb{R}^{4q}$. Note that the action φ commutes with A (see (17)) and both, A and φ, preserve the metric and are tri-holomorphic. Therefore, T^q also acts on the hyper-Kähler quotient by tri-holomorphic isometries. Moreover, the next theorem shows that when $l=p=q$, so that the quotient has dimension $4q$, the T^q-action has a unique fixed point.

In the next theorem we give the explicit description of the moment map and show that the quotient manifold is diffeomorphic to the Euclidean space.

Theorem 6.1. *Let $G_\theta = \mathbb{R}^s \times (\mathbb{R}^k \ltimes_\theta \mathbb{H}^q)$ be a hyper-Kähler Lie group, $s+k=4p$, $\theta \in M(k,q;k)$, L the connected closed abelian isotropic subgroup $L \subset \mathbb{R}^k$ defined above and A, φ as in* (17), (19). *Then*

1. *the expression of the moment map is*
$$\mu^\theta(X,W) = \left(-Im\, X_1 + \frac{1}{2}\sum_{\beta=1}^{q} \theta^1_\beta \overline{W_\beta}\, iW_\beta, \dots, -Im\, X_l + \frac{1}{2}\sum_{\beta=1}^{q} \theta^l_\beta \overline{W_\beta}\, iW_\beta \right), \tag{20}$$
for $(X,W) \in G_\theta$;
2. *we have the following diffeomorphisms:*
$$L\backslash(\mu^\theta)^{-1}(0) \cong \mathbb{R}^{4p+4q-4l}, \qquad L\backslash G_\theta \cong \mathbb{R}^{4p+4q-l};$$
3. *the torus T^q acts on $L\backslash(\mu^\theta)^{-1}(0)$ by tri-holomorphic isometries. If $l=p=q$, the action of T^q on the $4q$ dimensional quotient has a unique fixed point.*

Proof. In order to prove the second assertion we will find global coordinates on $L\backslash\mu_\theta^{-1}(0)$ and $L\backslash G_\theta$. For $(X,W)\in G_\theta$, $(T,0)\in L$, set

$$X = \sum_{\alpha=1}^{p} e_\alpha(x_\alpha + b_\alpha i + s_\alpha j + p_\alpha k), \qquad T = \sum_{\alpha=1}^{l} t_\alpha e_\alpha, \tag{21}$$

$$W = \sum_{\alpha=1}^{q} f_\alpha(u_\alpha + y_\alpha i + z_\alpha j + w_\alpha k). \tag{22}$$

It follows that $(x_\alpha, b_\gamma, s_\gamma, p_\gamma, u_\beta, y_\beta, z_\beta, w_\beta)$, with $\alpha = l+1,\dots,p$, $\gamma = l,\dots,p$, $\beta = 1,\dots,q$, are global coordinates on $L\backslash G_\theta$ and therefore $L\backslash G_\theta$ is diffeomorphic to $\mathbb{R}^{4p+4q-l}$. Using the fact that the hypercomplex structure corresponds to

$$J_1 = R_{-i}, \quad J_2 = R_{-j}, \quad J_3 = R_{-k}$$

and that the metric g is such that the real basis

$$\{e_\alpha, e_\alpha i, e_\alpha j, e_\alpha k, f_\beta, f_\beta i, f_\beta j, f_\beta k, \;\; 1\le\alpha\le p, \; 1\le\beta\le q\}$$

is orthonormal, we get the following expression of the moment maps ${\mu^\theta}_\gamma$, $\gamma = 1,2,3$, in terms of the real coordinates on $\mathbb{H}^p$ and $\mathbb{H}^q$:

$$\begin{aligned}
{\mu^\theta}_1(X,W)(T) &= -\sum_{\alpha=1}^{l} b_\alpha t_\alpha + \frac{1}{2}\sum_{\alpha=1}^{l} t_\alpha\left(\sum_{\beta=1}^{q}\theta^\alpha_\beta(u_\beta^2 + y_\beta^2 - z_\beta^2 - w_\beta^2)\right)\\
&= g\left(T, \sum_{\alpha=1}^{l}\left(-b_\alpha + \frac{1}{2}\sum_{\beta=1}^{q}\theta^\alpha_\beta(u_\beta^2 + y_\beta^2 - z_\beta^2 - w_\beta^2)\right)e_\alpha\right),\\
{\mu^\theta}_2(X,W)(T) &= -\sum_{\alpha=1}^{l} s_\alpha t_\alpha + \sum_{\alpha=1}^{l} t_\alpha\left(\sum_{\beta=1}^{q}\theta^\alpha_\beta(-u_\beta w_\beta + z_\beta y_\beta)\right)\\
&= g\left(T, \sum_{\alpha=1}^{l}\left(-s_\alpha + \sum_{\beta=1}^{q}\theta^\alpha_\beta(-u_\beta w_\beta + z_\beta y_\beta)\right)e_\alpha\right),\\
{\mu^\theta}_3(X,W)(T) &= -\sum_{\alpha=1}^{l} p_\alpha t_\alpha + \sum_{\alpha=1}^{l} t_\alpha\left(\sum_{\beta=1}^{q}\theta^\alpha_\beta(u_\beta z_\beta + w_\beta y_\beta)\right)\\
&= g\left(T, \sum_{\alpha=1}^{l}\left(-p_\alpha + \sum_{\beta=1}^{q}\theta^\alpha_\beta(u_\beta z_\beta + w_\beta y_\beta)\right)e_\alpha\right),
\end{aligned}$$

or, equivalently, (20) holds. On $(\mu^\theta)^{-1}(0)$ one has the following relations:

$$\begin{aligned} b_\alpha &= \frac{1}{2}\sum_{\beta=1}^{q}\theta^\alpha_\beta(u_\beta^2+y_\beta^2-z_\beta^2-w_\beta^2),\\ s_\alpha &= -\sum_{\beta=1}^{q}\theta^\alpha_\beta(u_\beta w_\beta - z_\beta y_\beta),\\ p_\alpha &= \sum_{\beta=1}^{q}\theta^\alpha_\beta(u_\beta z_\beta + w_\beta y_\beta),\end{aligned}$$

for any $\alpha = 1\ldots, l$. Thus, one has that $(x_\alpha, b_\gamma, s_\gamma, p_\gamma, u_\beta, y_\beta, z_\beta, w_\beta)$, with $\alpha = 1, \ldots, p$, $\gamma = l+1, \ldots, p$, $\beta = 1, \ldots, q$, are global coordinates on $(\mu^\theta)^{-1}(0)$.

Since the action of $\mathbb{R}^l$ leaves invariant $x_\gamma, b_\gamma, s_\gamma, p_\gamma$, $(\gamma \geq l+1)$ and rotates the coordinates $u_\beta, y_\beta, z_\beta, w_\beta$, $\beta = 1, \ldots, q$, one has that $(x_\gamma, b_\gamma, s_\gamma, p_\gamma, u_\beta, y_\beta, z_\beta, w_\beta)$, with $\gamma = l+1, \ldots, p$, $\beta = 1, \ldots, q$, are global coordinates on $L\backslash(\mu^\theta)^{-1}(0)$. It follows that the quotient space is diffeomorphic to $\mathbb{R}^{4p+4q-4l}$.

3. It follows from (19) that:

$$\varphi(B)(X, W) = (X, BW),$$

$B \in T^q, (X, W) \in G_\theta$. Since the moment map for the action φ is T^q-equivariant, φ preserves $(\mu^\theta)^{-1}(0)$. In particular, the hyper-Kähler quotient admits a tri-holomorphic action of the torus T^q.

Assume next that $l = p = q$, hence $\mathfrak{l} \oplus J_1\mathfrak{l} \oplus J_2\mathfrak{l} \oplus J_3\mathfrak{l} = \mathbb{R}^{4p}$. Let π be the natural projection from $(\mu^\theta)^{-1}(0)$ onto $L\backslash(\mu^\theta)^{-1}(0)$. Then $\pi(X, W)$, $(X, W) \in (\mu^\theta)^{-1}(0)$, is a fixed point for the action of T^q if and only if

$$(V, 0) \cdot (X, W) \cdot (V, 0)^{-1} \cdot (X, W)^{-1} \in L$$

for every $V \in \mathbb{R}^q$. We will show that $\pi(X, W) = \pi(0, 0)$, that is, $\pi(0, 0)$ is the unique fixed point. Using (19) and (12) we calculate

$$(V, 0) \cdot (X, W) \cdot (V, 0)^{-1} \cdot (X, W)^{-1} = (V, 0) \cdot (-V, W - (-V) \cdot W) = (0, V \cdot W - W)$$

which belongs to L if and only if $V \cdot W = W$ for every $V \in \mathbb{R}^q$, hence $W = 0$. Since $(X, 0) \in (\mu^\theta)^{-1}(0)$ it follows that $\omega_\alpha(V, X) = 0$ for every $V \in \mathfrak{l}$, $\alpha = 1, 2, 3$, and the assumption on l implies that $X \in \mathfrak{l}$. Therefore $\pi(X, 0) = \pi(0, 0)$, as asserted. □

Using the fact that the quotient admits a tri-holomorphic T^q-action, we can obtain the local expression of the hyper-Kähler metric in terms of the structure constants of the Lie group G_θ.

Observe that, if $l = p$, the quotient has dimension $4q$, thus by [10, 15] the induced hyper-Kähler metric can be locally written as follows

$$\frac{1}{4}H_{\beta\gamma}d\mathbf{r}_\beta \cdot d\mathbf{r}_\gamma + \frac{1}{4}H^{\beta\gamma}(d\tau_\beta + \mathbf{\Omega}_{\beta\delta} \cdot d\mathbf{r}_\delta)(d\tau_\gamma + \mathbf{\Omega}_{\gamma\epsilon} \cdot d\mathbf{r}_\epsilon), \tag{23}$$

where $\beta, \gamma = 1, \dots, q$, $(H)^{\beta\gamma}$ is the inverse of the matrix $(H)_{\beta\gamma}$. The Killing vector fields $\frac{\partial}{\partial \tau_\beta}$ generate the T^q-action, $\psi_\beta, \mathbf{r}_\beta$ are defined as in (15) and we assume Einstein summation convention. If $l < p$, the quotient splits as Riemannian product of the flat Euclidean space $\mathbb{R}^{4p-4l}$ by a $4q$-dimensional hyper-Kähler manifold with a tri-holomorphic T^q-action.

Theorem 6.2. *The local expression of the hyper-Kähler metric on the quotient* $L \backslash (\mu^\theta)^{-1}(0)$ *is* $h = h_0 + h_1$*, where* h_0 *is the Euclidean metric on* $\mathbb{R}^{4p-4l}$ *and* h_1 *is given by* (23)*, with*

$$H_{\beta\gamma} = (\tilde{\theta}\tilde{\theta}^t)_{\beta\gamma} + \frac{1}{r_\beta}\,\delta_{\beta\gamma}. \tag{24}$$

$\tilde{\theta}$ *is the* $q \times l$ *matrix obtained from* θ *by deleting the last* $p - l$ *columns,* $\tilde{\theta}^t$ *is its transpose and* $r_\beta = |\mathbf{r}_\beta|$.

Proof. The action A given by (17) of L on G_θ in the coordinates $(x_\alpha, b_\alpha, s_\alpha, p_\alpha, \psi_\beta, \mathbf{r}_\beta)$ is

$$\begin{aligned} &L \times G_\theta \to G_\theta, \\ &(T, (X_\alpha, \psi_\beta, \mathbf{r}_\beta)) \to (X_\alpha + t_\alpha, \psi_\beta + 2\langle \theta_\beta, T\rangle, \mathbf{r}_\beta), \end{aligned}$$

with $\alpha = 1, \dots, p$, $t_\alpha = 0$ for $\alpha > l$, $\beta = 1, \dots, q$ and $\psi_\beta, \mathbf{r}_\beta$ are defined as in (15).

The previous action leaves

$$\tau_\beta = \psi_\beta - 2 \sum_{\alpha=1}^{p} \theta_\beta^\alpha x_\alpha, \quad \beta = 1, \dots, q,$$

invariant and $\frac{\partial}{\partial \tau_\beta}$ are Killing vector fields for the quotient hyper-Kähler metric and generate the T^q-action induced by (19).

On $\mu^{-1}(0)$ one has

$$\mathrm{Im} X_\alpha = \frac{1}{2} \sum_{\beta=1}^{q} \theta_\beta^\alpha \mathbf{r}_\beta, \quad \alpha = 1, \dots, l,$$

so the metric on $\mu^{-1}(0)$ is

$$\sum_{\alpha=1}^{p} dx_\alpha^2 + \sum_{\alpha=l+1}^{p} d(\mathrm{Im} X_\alpha)^2 + \frac{1}{4} \sum_{\alpha=1}^{l} \left(\sum_{\beta=1}^{q} \theta_\beta^\alpha d\mathbf{r}_\beta \right)^2 + \frac{1}{4} \sum_{\beta=1}^{q} \left(\frac{1}{r_\beta} d\mathbf{r}_\beta^2 + r_\beta (d\psi_\beta + \mathbf{\Omega}_\beta \cdot d\mathbf{r}_\beta)^2 \right).$$

Projecting orthogonally to the Killing vector fields $\frac{\partial}{\partial x_\alpha}$, $\alpha = 1, \dots, l$, one gets that locally the metric on the quotient is given by $h = h_0 + h_1$, where

$$h_0 = \sum_{\alpha=l+1}^{p} (dx_\alpha^2 + d(\mathrm{Im} X_\alpha)^2)$$

and h_1 is given by (23) with the matrix H as in (24).

□

References

[1] D.V. Alekseevskii, B.N. Kimel'fel'd, *Structure of homogeneous Riemannian spaces with zero Ricci curvature*, Funktional Anal. i Proložen **9**(2), 5-11 (1975), (English translation: Functional Anal. Appl. **9**, 97-102 (1975)).

[2] R. Bielawski, A. S. Dancer, *The geometry and topology of toric hyper-Kähler manifolds*, Comm. Anal. Geom. 8 (2000), no. 4, 727-760.

[3] J. M. Dardie', A. Medina, *Algebres de Lie kähleriennes et double extension*, J. Algebra **185** (1996), no. 3, 774–795.

[4] G. W. Gibbons, P. Rychenkova, R. Goto, *Hyper-Kähler quotient construction of BPS monopole moduli spaces*, Comm. Math. Phys. **186** (1997), 581-599.

[5] R. Goto, *On toric hyper-Kähler manifolds given by the hyper-Kähler quotient method*, in Infinite analysis, Part A, B (Kyoto, 1991), 317-338, Adv. Ser. Math. Phys., 16, World Sci. Publishing, River Edge, NJ, 1992.

[6] V. Guillemin, S. Sternberg, *A normal form for the moment map*, in Differential geometric methods in mathematical physics, S.Sternberg editor (Reidel Publishing Company, 1984) pp. 161-175.

[7] R. Hermann, *A sufficient condition that a mapping of Riemannian manifolds be a fibre bundle*, Proc. A.M.S. **11**(1960), 236-242.

[8] N. J. Hitchin, A. Karlhede, U. Lindström, M. Roček, *Hyper-Kähler metrics and supersymmetry*, Comm. Math. Phys. **108** (1987), 535-589.

[9] O. Kowalski, F. Tricerri, L. Vanhecke, *Curvature homogeneous spaces with a solvable Lie group as homogeneous model*, J. Math. Soc. Japan **44** (1992), 461-484.

[10] U. Lindström, M. Roček, *Scalar tensor duality and* $N = 1, 2$ *nonlinear* σ*-models*, Nucl. Phys. B **222** (1983), 285-308.

[11] K. Lee, E. J. Weinberg, P. Yi, *The moduli space of many BPS monopoles for arbitrary gauge groups*, Phys.Rev. D **54** (1996) 1633-1643.

[12] J. Milnor, *Curvature of left-invariant metrics on Lie groups*, Adv. Math. **21** (1976), 293-329.

[13] M. K. Murray, *A note on the* $(1, 1, \ldots, 1)$ *monopole metric*, J. Geom. Phys. **23** (1997), No.1, 31-41.

[14] B. O'Neill, *The fundamental equations of a submersion*, Mich. Math. J. **13** (1966), 459-469.

[15] H. Pedersen, Y. S. Poon, *Hyper-Kähler metrics and a generalization of the Bogomolny equations*, Comm. Math. Phys. **117** (1988), 569-580.

[16] M. Roček, *Supersymmetry and nonlinear* σ*-models*, in Supersymmetry in physics (Los Alamos, N.M., 1983) Phys. D **15** (1985), 75-82.

Laura Barberis and Isabel G. Dotti: FaMAF, Universidad Nacional de Córdoba, 5000 Córdoba, Argentina

E-mail address: `barberis@mate.uncor.edu, idotti@mate.uncor.edu`

Anna Fino: Dipartimento di Matematica, Università di Torino, Via Carlo Alberto 10, 10123 Torino, Italy

E-mail address: `fino@dm.unito.it`